

Quark and lepton mass matrices with A_4 family symmetry ¹

Hideyuki Sawanaka ²

*Graduate School of Science and Technology, Niigata University, Ikarashi 2-8050, Niigata
950-2181, Japan*

Abstract

Realistic quark masses and mixing angles are obtained applying the successful A_4 family symmetry for leptons, motivated by the quark-lepton assignments of SU(5). The A_4 symmetry is suitable to give tri-bimaximal neutrino mixing matrix which is consistent with current experimental data. We study new scenario for the quark sector with the A_4 symmetry.

¹Talk given at International Workshop on Neutrino Masses and Mixings, University of Shizuoka, Shizuoka, Japan, 17-19 December 2006.

²e-mail address: hide@muse.sc.niigata-u.ac.jp

1 Introduction

The current observed neutrino mixing [1, 2] suggests around the maximal 2-3 mixing angle and zero 1-3 mixing angle: $\theta_{23} \sim \pi/4$, $\theta_{13} \sim 0$. In such a symmetric limit where both $\cos \theta_{23}$ and $|U_{e3}| = \sin \theta_{13}$ vanish, the resulting 3×3 effective Majorana neutrino mass matrix forms in the flavor basis as [3]

$$\begin{pmatrix} X & C & C \\ C & A & B \\ C & B & A \end{pmatrix}. \quad (1)$$

This matrix has an exact symmetric form under a Z_2 symmetry, i.e. the 2-3 (μ - τ) permutation, and is diagonalized by the unitary matrix:

$$U_{Z_2} = \begin{pmatrix} \cos \theta_{12} & -\sin \theta_{12} & 0 \\ \sin \theta_{12}/\sqrt{2} & \cos \theta_{12}/\sqrt{2} & -1/\sqrt{2} \\ \sin \theta_{12}/\sqrt{2} & \cos \theta_{12}/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}, \quad (2)$$

with remaining the solar mixing angle θ_{12} arbitrary and the entry X of (1) is determined as

$$X = A + B + \frac{2\sqrt{2}C}{\tan 2\theta_{12}}. \quad (3)$$

Now we well know two special values for θ_{12} which give typical mixing matrices; one is bimaximal and the other is tri-bimaximal mixing matrices. In a limit of bimaximal mixing [4] where $\theta_{12} = 4/\pi$, resulting MNS matrix forms

$$U_{\text{BM}} = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 1/2 & 1/2 & -1/\sqrt{2} \\ 1/2 & 1/2 & 1/\sqrt{2} \end{pmatrix}, \quad (4)$$

with

$$X = A + B. \quad (5)$$

For the case $\tan \theta_{12} = 1/\sqrt{2}$ with so-called tri-bimaximal mixing which is proposed by Harrison, Perkins and Scott, then we have the HPS type matrix [5, 6]:

$$U_{\text{HPS}} = \begin{pmatrix} 2/\sqrt{6} & 1/\sqrt{3} & 0 \\ -1/\sqrt{6} & 1/\sqrt{3} & -1/\sqrt{2} \\ -1/\sqrt{6} & 1/\sqrt{3} & 1/\sqrt{2} \end{pmatrix}, \quad (6)$$

where θ_{12} is fixed by as well

$$X = A + B + C \tag{7}$$

is also derived. Note that the tri-bimaximal structure is consistent with current experimental data, where θ_{12} is not maximal.

So far, the discrete symmetry A_4 is successfully applied for leptons. Namely, the tri-bimaximal mixing pattern can be realized naturally in a number of specific models [7, 8, 9, 10, 11]. However, it is not easy to have small quark mixing angles in naive and straight way. In such applications the generic prediction [8] is that V_{CKM} , the quark mixing matrix becomes just the unit matrix. Starting with $V_{\text{CKM}} = \mathbf{1}$, the realistic small quark mixing angles can be generated by extending interactions beyond those of the Standard Model, such as in supersymmetry [9] or breaking A_4 symmetry explicitly [10]. Our study is aimed to obtain realistic quark masses and mixing angles entirely within the A_4 context [12]. It is worthy of mention that the other types of unified models for quarks and leptons with the A_4 symmetry [13, 14] and models which predict tri-bimaximal mixing matrix with S_3 symmetry [15, 16, 17] have been also studied.

2 A_4 symmetry

A_4 is the symmetry group of the tetrahedron and the finite groups of the even permutation of four objects. It has twelve elements which are derived into four equivalence classes: $[C_1]$: (1234), $[C_2]$: (2143), (3412), (4321), $[C_3]$: (1342), (4213), (2431), (3124) and $[C_4]$: (1423), (3241), (4132), (2314), corresponding to its four irreducible representations we call three one-dimensional representations (singlets) as $\underline{1}$, $\underline{1}'$, $\underline{1}''$, and one three-dimensional representation (triplet) as $\underline{3}$, respectively. The A_4 is the smallest discrete group which includes the three-dimensional irreducible representation. The presence of the three-dimensional irreducible representation might be ideal for describing three families of quarks and leptons.

	h	n	$\underline{1}$	$\underline{1}'$	$\underline{1}''$	$\underline{3}$
C_1	1	1	1	1	1	3
C_2	2	3	1	1	1	-1
C_3	3	4	1	ω	ω^2	0
C_4	3	4	1	ω^2	ω	0

Table 1: Character table of A_4 .

The character table of four representations is shown in Table 1. Here h is the order of each element, n is the number of elements and the complex number ω is the cube root of unity:

$$\omega = \exp(2\pi i/3) = -\frac{1}{2} + \frac{\sqrt{3}}{2}i, \quad 1 + \omega + \omega^2 = 0. \quad (8)$$

The fundamental multiplication rules are given as³

$$\underline{1}' \times \underline{1}' = \underline{1}'', \quad \underline{1}'' \times \underline{1}'' = \underline{1}', \quad \underline{1}' \times \underline{1}'' = \underline{1}, \quad (9)$$

and

$$\underline{3}_1 \times \underline{3}_2 = \underline{1} + \underline{1}' + \underline{1}'' + \underline{3}_A + \underline{3}_B, \quad (10)$$

where denoting $\underline{3}_i$ for $i = 1, 2$ as (a_i, b_i, c_i) , we have

$$\begin{aligned} \underline{1} &\sim a_1 a_2 + b_1 b_2 + c_1 c_2, \\ \underline{1}' &\sim a_1 a_2 + \omega b_1 b_2 + \omega^2 c_1 c_2, \\ \underline{1}'' &\sim a_1 a_2 + \omega^2 b_1 b_2 + \omega c_1 c_2, \\ \underline{3}_A &\sim (b_1 c_2, c_1 a_2, a_1 b_2), \\ \underline{3}_B &\sim (c_1 b_2, a_1 c_2, b_1 a_2). \end{aligned}$$

Note that from Eq. (9) the A_4 invariant singlet $\underline{1}$ can be derived in these three sets of A_4 singlets:

$$\underline{1}' \times \underline{1}' \times \underline{1}', \quad \underline{1}'' \times \underline{1}'' \times \underline{1}'', \quad \underline{1} \times \underline{1}' \times \underline{1}'', \quad (11)$$

³For details of the A_4 multiplication rules, see the original paper [7, 8] for example.

	(ν_i, l_i)	e^c	μ^c	τ^c	ϕ_{li}
A_4	$\underline{\mathbf{3}}$	$\underline{\mathbf{1}}$	$\underline{\mathbf{1}}'$	$\underline{\mathbf{1}}''$	$\underline{\mathbf{3}}$

Table 2: A_4 assignment for leptons.

and $\underline{\mathbf{3}} \times \underline{\mathbf{3}} \times \underline{\mathbf{3}} = \underline{\mathbf{1}}$ is also possible in the A_4 symmetry from Eq. (10). By using them, A_4 invariant mass matrices are constructed.

3 A_4 model for lepton sector

In this section, we briefly show a simple example of the A_4 model for leptons and lead the tri-bimaximal mixing matrix. For more detail of advanced models can be found in recent reviews [18, 19].

Let us take the A_4 assignment for leptons as shown in Table 2: left-handed $SU(2)_L$ lepton doublets (ν_i, l_i) ($i = 1, 2, 3$) transform as an A_4 triplet $\underline{\mathbf{3}}$, while right-handed, charged lepton singlets l_i^c transform as A_4 singlets ($l_1^c = e^c$, $l_2^c = \mu^c$ and $l_3^c = \tau^c$ transform as $\underline{\mathbf{1}}$, $\underline{\mathbf{1}}'$ and $\underline{\mathbf{1}}''$, respectively). Introducing gauge singlet Higgs doublet $\phi_{li} = (\phi_{li}^0, \phi_{li}^-) \sim \underline{\mathbf{3}}$ under A_4 , the 3×3 mass matrix linking l_i with l_i^c is given by

$$M_l = \begin{pmatrix} f_1 v_{l1} & f_2 v_{l1} & f_3 v_{l1} \\ f_1 v_{l2} & f_2 \omega v_{l2} & f_3 \omega^2 v_{l2} \\ f_1 v_{l3} & f_2 \omega^2 v_{l3} & f_3 \omega v_{l3} \end{pmatrix} = \begin{pmatrix} v_{l1} & 0 & 0 \\ 0 & v_{l2} & 0 \\ 0 & 0 & v_{l3} \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} f_1 & 0 & 0 \\ 0 & f_2 & 0 \\ 0 & 0 & f_3 \end{pmatrix}, \quad (12)$$

where $\omega = \exp(2\pi i/3)$, f_i are Yukawa couplings and $v_{li} = \langle \phi_{li}^0 \rangle$ are vacuum expectation values of the Higgs field ϕ_{li}^0 . It can be diagonalized in a very simple way, i.e. by setting all three vacuum expectation values of the Higgs field to be equal. Taking $v_l \equiv v_{l1} = v_{l2} = v_{l3}$, the charged lepton mass matrix is diagonalized as

$$M_l^{diag} = U_{lL}^\dagger M_l U_{lR} = \begin{pmatrix} f_1 & 0 & 0 \\ 0 & f_2 & 0 \\ 0 & 0 & f_3 \end{pmatrix} \sqrt{3} v_l = \begin{pmatrix} m_e & 0 & 0 \\ 0 & m_\mu & 0 \\ 0 & 0 & m_\tau \end{pmatrix}, \quad (13)$$

where

$$U_{lL} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}, \quad (14)$$

and U_{lR} is the unit matrix. Three different charged lepton masses are given by the Yukawa couplings.

For the neutrino mass matrix, if we consider six gauge singlet Higgs triplets $\xi_i = (\xi_i^{++}, \xi_i^+, \xi_i^0)$ which are assigned to $\xi_1 \sim \underline{1}$, $\xi_2 \sim \underline{1}'$, $\xi_3 \sim \underline{1}''$ and $\xi_{4,5,6} \sim \underline{3}$ under A_4 , then the matrix forms in general

$$M_\nu = \begin{pmatrix} a+b+c & f & e \\ f & a+\omega b+\omega^2 c & d \\ e & d & a+\omega^2 b+\omega c \end{pmatrix}, \quad (15)$$

here parameters (a, b, c) come from (ξ_1, ξ_2, ξ_3) and (d, e, f) from $\xi_{4,5,6}$, respectively. If conditions $b = c$ and $e = f = 0$ are given (hopefully in some natural mechanisms), M_ν is diagonalized by the matrix $U_{\nu L}$ with three eigenvalues (neutrino masses) $m_1 = a - b + d$, $m_2 = a + 2b$, $m_3 = -a + b + d$, and then we have

$$U_{\text{MNS}} = (U_{lL})^\dagger U_{\nu L} = \begin{pmatrix} 2/\sqrt{6} & 1/\sqrt{3} & 0 \\ -1/\sqrt{6} & 1/\sqrt{3} & -1/\sqrt{2} \\ -1/\sqrt{6} & 1/\sqrt{3} & 1/\sqrt{2} \end{pmatrix}. \quad (16)$$

which is exactly the tri-bimaximal mixing matrix.

4 A_4 model for quark sector

Following the successful lepton assignments in the previous section, if we assign for quarks as in Table 3 and take $v_{q1} = v_{q2} = v_{q3} \equiv v_q$ ($v_{qi} = \langle \phi_{qi}^0 \rangle$), we then have the up and down quark mass matrices as

$$M^{U(D)} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} g_1^{U(D)} & 0 & 0 \\ 0 & g_2^{U(D)} & 0 \\ 0 & 0 & g_3^{U(D)} \end{pmatrix} \sqrt{3} v_q. \quad (17)$$

	(u_i, d_i)	u_i^c, d_i^c	$(\phi_{qi}^0, \phi_{qi}^-)$
A_4	$\underline{\mathbf{3}}$	$\underline{\mathbf{1}}, \underline{\mathbf{1}}', \underline{\mathbf{1}}''$	$\underline{\mathbf{3}}$

Table 3: A_4 assignment for quarks which follows that for leptons.

	$(\nu_i, l_i), d_i^c$	$l_i^c, u_i^c, (u_i, d_i)$	$(\phi_{U1,2}^0, \phi_{U1,2}^-)$	$(\phi_{li}^0, \phi_{li}^-), (\phi_{Di}^0, \phi_{Di}^-)$
A_4	$\underline{\mathbf{3}}$	$\underline{\mathbf{1}}, \underline{\mathbf{1}}', \underline{\mathbf{1}}''$	$\underline{\mathbf{1}}', \underline{\mathbf{1}}''$	$\underline{\mathbf{3}}$
SU(5)	$\underline{\mathbf{5}}^*$	$\underline{\mathbf{10}}$	$\underline{\mathbf{5}}$	$\underline{\mathbf{5}}^* + \underline{\mathbf{45}}$

Table 4: SU(5) motivated quark and lepton assignment.

The generic prediction is that the quark mixing matrix leads to the unit matrix: $V_{\text{CKM}} = (U^U)^\dagger U^D = \mathbf{1}$ where

$$U^U = U^D = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \quad (18)$$

are matrices which diagonalize $M^{U(D)}$ as (13). The realistic small quark mixing angles can be generated by extending interactions beyond those of the Standard Model, such as in supersymmetry [10] or the addition of terms which break the A_4 symmetry as well as the residual Z_3 symmetry explicitly [11]. In this section, we present a new alternative scenario, where realistic quark masses and mixing angles are obtained motivated by the quark-lepton assignments of SU(5), entirely within the A_4 context [12].

In SU(5) grand unification, the $\underline{\mathbf{5}}^*$ representation contains the lepton doublet (ν, l) and the quark singlet d^c , whereas the $\underline{\mathbf{10}}$ representation contains the lepton singlet l^c and the quark doublet (u, d) and singlet u^c . In the successful A_4 model for leptons, (ν_i, l_i) transform as $\underline{\mathbf{3}}$ whereas l_i^c transform as $\underline{\mathbf{1}}, \underline{\mathbf{1}}'$ and $\underline{\mathbf{1}}''$. Thus we choose as shown in Table 4. In minimal SU(5), there is just one $\underline{\mathbf{5}}$ representation of Higgs bosons, yielding thus only two invariants, i.e. $\underline{\mathbf{10}} \times \underline{\mathbf{10}} \times \underline{\mathbf{5}} \rightarrow \underline{\mathbf{1}}$ (that is for up quark mass matrix) and $\underline{\mathbf{5}}^* \times \underline{\mathbf{10}} \times \underline{\mathbf{5}}^* \rightarrow \underline{\mathbf{1}}$ (for charged lepton and down quark mass matrices). The second invariant implies $m_\tau = m_b$ at the unification

scale which is phenomenologically desirable. We follow the usual strategy of using both $\underline{5}^*$ and $\underline{45}$ representations of Higgs bosons, so that one linear combination couples to only leptons, and the other only to quarks. Both transform as $\underline{3}$ under A_4 . There are also two $\underline{5}$ representations transforming as $\underline{1}'$ and $\underline{1}''$ under A_4 which couple only to up type quarks⁴.

With this A_4 assignment for Higgs doublets, the relevant Yukawa couplings linking d_i with d_j^c are given by

$$\begin{aligned} & h_1 d_1 (d_1^c \phi_{D1}^0 + d_2^c \phi_{D2}^0 + d_3^c \phi_{D3}^0) + h_2 d_2 (d_1^c \phi_{D1}^0 + \omega d_2^c \phi_{D2}^0 + \omega^2 d_3^c \phi_{D3}^0) \\ & + h_3 d_3 (d_1^c \phi_{D1}^0 + \omega^2 d_2^c \phi_{D2}^0 + \omega d_3^c \phi_{D3}^0), \end{aligned}$$

resulting in the 3×3 down quark mass matrix:

$$M_D = \begin{pmatrix} h_1 & 0 & 0 \\ 0 & h_2 & 0 \\ 0 & 0 & h_3 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} v_1 & 0 & 0 \\ 0 & v_2 & 0 \\ 0 & 0 & v_3 \end{pmatrix}, \quad (19)$$

where $\omega = \exp(2\pi i/3)$, h_i are three independent Yukawa couplings, and v_i are the vacuum expectation values of ϕ_{Di}^0 . To get quark mass hierarchy, $v_1 \ll v_2 \ll v_3$ should be satisfied contrary to the lepton sector ($v_{li} \equiv \langle \phi_{li}^0 \rangle = v_l$). By contrast, the Higgs doublets transforming as $\underline{1}'$ and $\underline{1}''$ linking u_i with u_j^c give, as following the manner of Eq. (11), the 3×3 symmetric up quark mass matrix:

$$M_U = \begin{pmatrix} 0 & \mu_2 & \mu_3 \\ \mu_2 & m_2 & 0 \\ \mu_3 & 0 & m_3 \end{pmatrix}, \quad (20)$$

where m_2, μ_3 come from $\phi_{U1}^0 \sim \underline{1}'$ and m_3, μ_2 from $\phi_{U2}^0 \sim \underline{1}''$ and three of them can be taken real in general. In the limit $|\mu_2| \ll |m_2|$ and $|\mu_3| \ll |m_3|$, we obtain the three eigenvalues of M_U as

$$m_t \simeq |m_3|, \quad m_c \simeq |m_2|, \quad m_u \simeq \left| \frac{\mu_2^2}{m_2} + \frac{\mu_3^2}{m_3} \right|, \quad (21)$$

⁴We choose here two Higgs doublets transforming as $\underline{1}'$ and $\underline{1}''$, other patterns can be assigned with different prediction.

with mixing angles

$$V_{uc} \simeq \frac{\mu_2}{m_2}, \quad V_{ut} \simeq \frac{\mu_3}{m_3}, \quad V_{ct} \simeq 0. \quad (22)$$

In the down sector, we note that

$$M_D M_D^\dagger = \begin{pmatrix} Y|h_1|^2 & Z^*h_1h_2^* & Zh_1h_3^* \\ Zh_1^*h_2 & Y|h_2|^2 & Z^*h_2h_3^* \\ Z^*h_1^*h_3 & Zh_2^*h_3 & Y|h_3|^2 \end{pmatrix}, \quad (23)$$

where

$$\begin{aligned} Y &= |v_1|^2 + |v_2|^2 + |v_3|^2, \\ Z &= |v_1|^2 + \omega|v_2|^2 + \omega^2|v_3|^2. \end{aligned}$$

Its eigenvalue λ satisfies the equation

$$\begin{aligned} &\lambda^3 - Y(|h_1|^2 + |h_2|^2 + |h_3|^2)\lambda^2 - (Y^3 + Z^3 + Z^{*3} - 3Y|Z|^2)|h_1|^2|h_2|^2|h_3|^2 \\ &+ (Y - |Z|^2)(|h_1|^2|h_2|^2 + |h_1|^2|h_3|^2 + |h_2|^2|h_3|^2)\lambda = 0. \end{aligned} \quad (24)$$

If $|v_1| = |v_2| = |v_3| = |v|$ as well as assumed in the charged lepton case, then $Y = 3|v|^2$, $Z = 0$ and three eigenvalues are simply $3|h_{1,2,3}|^2|v|^2$. We choose them instead to be different, but we still assume $|h_1|^2 \ll |h_2|^2 \ll |h_3|^2$. In that case, we find

$$m_b^2 \simeq Y|h_3|^2, \quad (25)$$

$$m_s^2 \simeq \left(\frac{Y^2 - |Z|^2}{Y} \right) |h_2|^2, \quad (26)$$

$$m_d^2 \simeq \left(\frac{Y^3 + Z^3 + Z^{*3} - 3Y|Z|^2}{Y^2 - |Z|^2} \right) |h_1|^2, \quad (27)$$

and the mixing angles are given by

$$V_{sb} \simeq \left(\frac{Z^*}{Y} \right) \frac{h_2}{h_3}, \quad (28)$$

$$V_{db} \simeq \left(\frac{Z}{Y} \right) \frac{h_1}{h_3}, \quad (29)$$

$$V_{ds} \simeq \left(\frac{YZ^* - Z^2}{Y^2 - |Z|^2} \right) \frac{h_1}{h_2}, \quad (30)$$

thereby requiring the condition

$$\left| \frac{V_{ds}V_{sb}}{V_{db}} \right| \simeq \left| \frac{YZ^* - Z^2}{Y^2 - |Z|^2} \right|. \quad (31)$$

Using current experimental values for the left-hand side, we see that quark mixing in the down sector alone cannot explain the observed quark mixing matrix V_{CKM} . Taking into account V_U , we then have

$$V_{\text{CKM}} = V_U^\dagger V_D. \quad (32)$$

Hence

$$V_{us} \simeq V_{ds} - V_{uc} \simeq \left(\frac{YZ^* - Z^2}{Y^2 - |Z|^2} \right) \frac{h_1}{h_2} - \frac{\mu_2}{m_2}, \quad (33)$$

$$V_{cb} \simeq V_{sb} \simeq \left(\frac{Z^*}{Y} \right) \frac{h_2}{h_3}, \quad (34)$$

$$V_{ub} \simeq V_{db} - V_{uc}V_{sb} - V_{ut} \simeq \left(\frac{Z}{Y} \right) \frac{h_1}{h_3} - \left(\frac{Z^*}{Y} \right) \frac{h_2}{h_3} \frac{\mu_2}{m_2} - \frac{\mu_3}{m_3}. \quad (35)$$

It is noted that our up and down quark mass matrices are restricted by our choice of A_4 representations to have only five independent parameters each. In the up sector, we have three real and one complex parameters (for example we choose here m_2 , m_3 and μ_2 to be real with μ_3 complex). The five independent parameters can be chosen as the three up quark masses, one mixing angle and one phase. In the down sector, the Yukawa couplings $h_{1,2,3}$ can all be chosen real, Y is just an overall scale, and Z is complex. The five independent parameters can be chosen as the three down quark masses and two mixing angles. Now we have ten parameters in these two matrices except their overall normalizations or magnitudes of Yukawa couplings. Since we also have ten observables for six quark masses, three angles and one phase, it may appear that a fit is not so remarkable. However, the forms of the mass matrices are very restrictive, and it is by no means trivial to obtain a good fit. Indeed, we find that V_{ub} is strongly correlated with the CP phase β which is one of angles of the unitary triangle. If we were to fit just the six masses and the three angles, the structure of our mass

matrices would allow only a very narrow range of values for β at each value of $|V_{ub}|$. This means that future more precise determinations of these two parameters will be a decisive test of this model.

CP violation is also predicted in our model. The Jarlskog invariant [23] is given by

$$J_{\text{CP}} \simeq \frac{\sqrt{3}}{2} \frac{h_1^2}{h_3^2} \left(\frac{v_2^2 - v_1^2}{v_2^2 + v_1^2} \right) \left(1 + \frac{\text{Re}(\mu_3) + \frac{1}{\sqrt{3}}\text{Im}(\mu_3) \frac{h_3}{h_1}}{m_t} \right), \quad (36)$$

which is mainly comes from the down sector and the up sector only affects as correction terms.

In order to fit the ten observables (six quark masses, three CKM mixing angles and one CP phase), ten parameters of our model have been generated numerically. We choose the parameter sets which are allowed by the experimental data. We show the prediction of $|V_{ub}|$ verses β in Fig. 1, with the following nine experimental inputs [20, 21, 22]:

$$\begin{aligned} m_u &= 0.9 \sim 2.9 \text{ (MeV)}, & m_c &= 530 \sim 680 \text{ (MeV)}, & m_t &= 168 \sim 180 \text{ (GeV)}, \\ m_d &= 1.8 \sim 5.3 \text{ (MeV)}, & m_s &= 35 \sim 100 \text{ (MeV)}, & m_b &= 2.8 \sim 3 \text{ (GeV)}, \\ |V_{us}| &= 0.221 \sim 0.227, & |V_{cb}| &= 0.039 \sim 0.044, & J_{\text{CP}} &= (2.75 \sim 3.35) \times 10^{-5}, \end{aligned} \quad (37)$$

which are given at the electroweak scale. We see that the experimental allowed region of β ($0.370 \sim 0.427$ radian at 90% C.L.) [22] corresponds to $|V_{ub}|$ in the range $0.0032 \sim 0.0044$, which is consistent with the experimental value of $|V_{ub}| = 0.0029 \sim 0.0045$. Thus our model is able to reproduce realistically the experimental data of quark masses and the CKM matrix.

Precisely measured heavy quark masses and CKM matrix elements are expected in future experiments and precise light quark masses are expected in future lattice evaluations. If the allowed regions of the current data shown in Eq. (37) are reduced, the correlation between $|V_{ub}|$ and β will become stronger. We show in Fig. 2 the case where the experimental data




Figure 1: Plot of allowed values in the $\beta - |V_{ub}|$ plane, where the value of β is expressed in radians. The horizontal and vertical lines denote experimental bounds at 90% C.L. (37).

are restricted to some very narrow ranges about their central values:

$$\begin{aligned}
 m_u &= 1.4 \sim 1.5 \text{ (MeV)}, & m_c &= 600 \sim 610 \text{ (MeV)}, & m_t &= 172 \sim 176 \text{ (GeV)}, \\
 m_d &= 3.4 \sim 3.6 \text{ (MeV)}, & m_s &= 60 \sim 70 \text{ (MeV)}, & m_b &= 2.85 \sim 2.95 \text{ (GeV)}, \\
 |V_{us}| &= 0.221 \sim 0.227, & |V_{cb}| &= 0.041 \sim 0.042, & J_{\text{CP}} &= (3.0 \sim 3.1) \times 10^{-5},
 \end{aligned}
 \tag{38}$$

Here we use the tighter constraints on the mass ratios of light quarks, i.e. m_u/m_d and m_s/m_d , consistent with the well-known successful low-energy sum rules [24]: Clearly, future more precise determinations of $|V_{ub}|$ and β will be a sensitive test of our model.

A comment is in order. Our quark mass matrices are in principle given at the SU(5) unification scale. However, the A_4 flavor symmetry is spontaneously broken at the electroweak scale. Therefore, the forms of our mass matrices are not changed except for the magnitudes of the Yukawa couplings between the unification and electroweak scales. Hence,




Figure 2: Plot of allowed values in the $\beta - |V_{ub}|$ plane, where input data are restricted in the narrower regions shown in Eq (38).

our numerical analyses are presented at the electroweak scale.

We should also comment on the hierarchy of h_i and v_i . The order of h_i are fixed by the quark mixing (33), (34). The ratios of

$$h_1/h_2 \simeq \lambda (\simeq 0.22), \quad h_2/h_3 \simeq \lambda^2, \quad (39)$$

are required by V_{us} and V_{cb} , respectively. Once h_i are fixed, quark masses determine the hierarchy of v_i as follows:

$$v_1/v_3 \simeq \lambda^2, \quad v_2/v_3 \simeq \lambda \sim \lambda^{1/2}. \quad (40)$$

These hierarchies of h_i and v_i are also consistent with the magnitude of J_{CP} given in Eq. (36).

5 Summary

The A_4 family symmetry which has been successful in understanding the mixing pattern of neutrinos (tri-bimaximal mixing) is applied to quarks, motivated by the quark-lepton

assignments of $SU(5)$. Realistic quark masses and mixing angles are obtained entirely with the A_4 context, in good agreement with data. In particular, we find a strong correlation between V_{ub} and the CP phase β , thus a decisive future test of this model can be allowed.

It is one of a powerful guideline to find the constraints from neutrinos to grand unification models. Discrete symmetries are suitable to decode the flavor problem: they can accommodate maximal 2-3 mixing and explain zero 1-3 mixing. Moreover they can be the origin of texture zeros or equalities.

Acknowledgements

I would like to thank Y. Koide for his great hospitality and organization of this stimulating workshop in Shizuoka. I am also grateful to E. Ma and M. Tanimoto for their collaboration.

References

- [1] M. Maltoni, T. Schwetz, M.A. Tortola and J.W.F. Valle, *New J. Phys.* **6** (2004) 122.
- [2] G.L. Fogli, E. Lisi, A. Marrone and A. Palazzo, *Prog. Part. Nucl. Phys.* **57** (2006) 742.
- [3] See for example, W. Grimus, A.S. Joshipura, S. Kaneko, L. Lavoura, H. Sawanaka and M. Tanimoto, *Nucl. Phys. B* **713** (2005) 151, and references therein.
- [4] V.D. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, *Phys. Lett. B* **437** (1998) 107.
- [5] P.F. Harrison, D.H. Perkins and W.G. Scott, *Phys. Lett. B* **530** (2002) 167; P.F. Harrison and W.G. Scott, *Phys. Lett. B* **535** (2002) 163.
- [6] See also, Z.Z. Xing, *Phys. Lett. B* **533** (2002) 85; X.G. He and A. Zee, *Phys. Lett. B* **560** (2003) 87.

- [7] E. Ma and G. Rajasekaran, Phys. Rev. D **64** (2001) 113012.
- [8] K.S. Babu, E. Ma and J.W.F. Valle, Phys. Lett. B **552** (2003) 207.
- [9] K.S. Babu, B. Dutta and R.N. Mohapatra, Phys. Rev. D **60** (1999) 095004.
- [10] E. Ma, Mod. Phys. Lett. A **17** (2002) 627; X.G. He, Y.Y. Keum and R.R. Volkas, JHEP **0604** (2006) 039.
- [11] See for example, M. Hirsch, J.C. Romao, S. Skadhauge, J.W.F. Valle and A. Villanova del Moral, Phys. Rev. D **69** (2004) 093006; E. Ma, Phys. Rev. D **70** (2004) 031901R; Phys. Rev. D **72** (2005) 037301; Mod. Phys. Lett. A **20** (2005) 2601; Phys. Rev. D **73** (2006) 057304; G. Altarelli and F. Feruglio, Nucl. Phys. B **720** (2005) 64; Nucl. Phys. B **741** (2006) 215; Nucl. Phys. B **724** (2005) 423; K.S. Babu and X.G. He, hep-ph/0507217; A. Zee, Phys. Lett. B **630** (2005) 58; B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma and M.K. Parida, Phys. Lett. B **638** (2006) 345; L. Lavoura and H. Kuhbock, Mod. Phys. Lett. A **22** (2007) 181; G. Altarelli, F. Feruglio and Y. Lin, hep-ph/0610165.
- [12] E. Ma, H. Sawanaka and M. Tanimoto, Phys. Lett. B **641** (2006) 301.
- [13] E. Ma, Mod. Phys. Lett. A **21** (2006) 2931.
- [14] S.F. King and M. Malinsky, Phys. Lett. B **645** (2007) 351.
- [15] W. Grimus and L. Lavoura, JHEP **0601** (2006) 018.
- [16] N. Haba, A. Watanabe and K. Yoshioka, Phys. Rev. Lett. **97** (2006) 041601.
- [17] R.N. Mohapatra, S. Nasri and H.B. Yu, Phys. Lett. B **639** (2006) 318.
- [18] R.N. Mohapatra and A.Y. Smirnov, Ann. Rev. Nucl. Part. Sci. **56** (2006) 569.
- [19] G. Altarelli, hep-ph/0611117.

- [20] H. Fritzsch and Z.Z. Xing, Prog. Part. Nucl. Phys. **45** (2000) 1.
- [21] Particle Data Group, W.M. Yao *et al.*, J. Phys. G **33** (2006) 1.
- [22] CKM fitter group, J. Charles, hep-ph/0606046.
- [23] C. Jarlskog, Phys. Rev. Lett. **55** (1985) 1039.
- [24] H. Leutwyler, Phys. Lett. B **378** (1996) 313; hep-ph/9609467.